

Applying Tags

Tags reflect the qualitative content of underlying studies and provide method for attaching text or images from these studies. After tags have been [configured](#), and so long as at least one study has been included, you can begin applying tags. Once a tag is applied, it is immediately viewable on [Qualitative Synthesis](#).

Since nests are set to Form-based Tagging mode by default, if you have configured Questions, follow [this link to learn how to use Form-based Tagging specifically](#). If not, the standard “Tagging” tab with the dropdown is still available underneath questions on the right handside. If you prefer to only see this tab and not the Questions tab, switch to Standard Tagging (Settings -> Tagging -> toggle Standard).



Regardless of Tagging mode, the below functions are both applicable. We would recommend going through this page and the above Form-based-specific page for guidance on applying tags/collecting data.

Steps for Tagging:

1. Navigate to Tagging

Click the “Tagging” button on the left-hand side, in the Nest Menu.

The screenshot displays the Nested Knowledge interface for a protocol titled "Home: Thrombectomy alone vs. Thrombectomy plus thrombolysis". On the left, the "Nest Menu" is visible, with the "Tagging" option highlighted. The main content area shows the "Protocol" details, including the Objective, Scope, Population, Primary Outcomes, Secondary Outcomes, and Interventions and Comparators. The right sidebar shows a chat window with a message from Kevin Kallmes.

This will enable you to apply tags to records sequentially. If you would prefer to search and find

records to tag, or to view records that have already been tagged, use [Study Inspector](#).

2. View the Full Text

Click on the “Full Text” toggle in the upper left to view the full-text PDF.

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Nest Home

Dashboard

Settings

Literature Search

1 / 1

Other Sources

Duplicate Review

Search Exploration

Query Builder

Screening

271 / 274

Configure Screening

Tagging

3 / 6

Configure Tagging

Extraction

6 / 6

Configure Extraction

Risk of Bias

6 / 6

Study Inspector

Synthesis

Manuscript Editor

Abstract Editor

Export

Abstract

Full Text

Supplements

Related Reports

10

0

4

0

PubMed

Navigation

Back

Skip

Complete

Tagging

Tag

Text

Select Tag

Enter Text

Apply Tag

Tag Recommendations

Comments (0)

History

Articles

Endovascular thrombectomy versus standard bridging thrombolytic with endovascular thrombectomy within 4-5 h of stroke onset: an open-label, blinded-endpoint, randomised non-inferiority trial

Peter J Mitchell*, Bernard Yan*, Leonid Churilov, Richard J Dowling, Steven J Bush, Andrew Bivard, Xiao Chuan Huo, Guoqing Wang, Shi Yong Zhang, Mai Duy Ton, Dennis J Cordato, Timothy J Kleing, Henry Ma, Ronil V Chandra, Helen Brown, Bruce CV Campbell, Andrew K Cheung, Brendan Steinfert, Rebecca Scroop, Kendal Redmond, Ferdinand Mikeff, Yan Liu, Dang Phuc Duc, Hal Rice, Mark W Parsons, Teddy Y Wu, Huy-Thang Nguyen, Geoffrey A Donnan†, Zhong Rong Miao†, Stephen M Davis†, on behalf of the DIRECT-SAFE Investigators†

Lancet 2022; 400: 116–25

See Comment page 74

*Joint principal investigators

†Joint senior authors

Investigators listed at the end of the report

Department of Radiology (Prof P J Mitchell MBBS, R J Dowling MBBS, S J Bush MBBS), Department of Medicine and Neurology, Melbourne Brain Centre (Prof B Yan DMScSc), A Bivard PhD, Prof B CV Campbell PhD, Prof G A Donnan MD,

If no full text has yet been imported, learn how to upload it [here](#).

3. Find the Relevant Tag

As you read through the article and find the relevant tags in the text, tables, or figures, search/select the relevant tag in the drop-down:



Tags are ordered in the dropdown based on the hierarchy with the leftmost root node at the top, followed by its children, followed by the next root node.

3a. Tag Details

If you need further details on the tags in order to determine applicability, and if the Tag Description was filled in for the tag in question, you can view it next to the Tag drop-down. An “i” icon will appear next to the Tag if a Description exists, and you can view it upon hovering:



3b. Tag Recommendations

Regardless of your Tagging mode, there will be a right-hand menu tab titled “Tag Recommendations”

beneath either “Tagging” (in Standard mode) or “Questions” (in Form-based mode). Tag Recommendations searches the study full text, highlighting specific text that may be applicable to the tags in your hierarchy. Standard Tag Recommendations (available to all users) perform a key word search of the tag name, while Smart Tag Recommendations (only available to enterprise users) utilize OpenAI GPT 3.5/4 to perform a smart search of the tag data. Standard tags are automatic, while Smart Tags can be switched on in Settings and used to generate recommendations for Abstracts as well as Full Texts.

Learn more about how to use [Tag Recommendations](#).

4. Add an Annotation

To associate text content with a tag, identify this text either before or after selecting the tag from the drop-down. You have three options for how to identify the text excerpt that will be associated with that tag:

- **Highlighting (Text Annotation):** A traceable, exact quote from the text of the article.
- **Selection (Area Annotation):** A traceable, exact image extraction from a table, figure, or other area of the article.
- **Manual entry (No Annotation):** A non-traceable excerpt (that is, an excerpt that is not connected to a specific part of the article) that you type into the Tag Text box.

You can annotate either before or after selecting the tag of interest in the drop-down (red circle below).

4a. Use the Highlighting Tool:

The screenshot displays the 'Tagging' interface for the article 'Endovascular thrombectomy versus standard bridging thrombolytic with endovascular thrombectomy within 4-5 h of stroke onset: an open-label, blinded-endpoint, randomised non-inferiority trial'. The interface includes a left sidebar with navigation options like 'Nest Home', 'Literature Search', 'Screening', 'Tagging', 'Extraction', 'Risk of Bias', 'Study Inspector', and 'Synthesis'. The main content area shows the article title and abstract. A red circle highlights the 'Highlighting' icon in the top toolbar. On the right, a 'Navigation' panel shows the 'Tagging' section with a dropdown menu set to 'Onset-to-needle' and a text input field for 'Enter Text'. Below this, there are sections for 'Tag Recommendations', 'Comments (0)', and 'History'.

The default Tag Text method is Highlighting. You can also manually select the Highlighting icon (see red arrow above), if you need to toggle back to this option.

Click and drag over the text you would like to Highlight. Highlighting will extract an exact text excerpt that is shown in light blue (see red arrow below), and the text will be automatically populated to the Tag Text box (red outline below).

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Nest Home

Literature Search

Screening

Tagging

Extraction

Risk of Bias

Study Inspector

Synthesis

Abstract

Full Text

Supplements

Related Reports

10

0

0

4

0

0

PubMed

Navigation

Tagging

Tag Recommendations

Comments (0)

History

Thrombectomy mTICI score ≥3

127/143 (89%)

130/146 (89%)

Adjusted OR 0.54 (0.39 to 0.82)

p=0.66

NIHSS score within 72 h

4 (1-11), n=141

4 (1-11), n=142

Adjusted OR 0.73 (0.45 to 1.18)

p=0.20

Early neurological improvement*

84/141 (60%)

95/142 (67%)

Adjusted OR 0.73 (0.45 to 1.18)

p=0.20

Safety outcomes

Death

22/146 (15%)

24/147 (16%)

Adjusted OR 0.92 (0.46 to 1.84)

p=0.82

Symptomatic intracerebral haemorrhage

2/146 (1%)

3/147 (2%)

Adjusted OR 1.70 (0.22 to 13.04)

p=0.62

Any intracerebral haemorrhage

3/146 (2%)

3/147 (2%)

Adjusted OR 0.97 (0.56 to 1.70)

p=0.92

Data are n/N (%) or median (IQR). ITT=Intention to treat; mRS=modified Rankin Scale; PP=per protocol; OR=odds ratio; NIHSS=National Institutes of Health Stroke Scale; mTICI=modified Treatment in Cerebral Ischemia; *NIHSS reduction of 8 points or more, or reaching 0-1 at 3 days, adjusted for baseline NIHSS and age.

Table 3: Study outcomes

patients lost to follow-up. In the per-protocol analysis

times from arterial puncture to reperfusion were below

after patient crossover, 145 remained in the direct

60 min in both groups (55-5 min [26-0-88-5] in the direct

thrombectomy group and 143 in the bridging therapy

thrombectomy group and 44-5 min [27-0-70-0] in the

group (figure 1). In the direct thrombectomy group,

bridging group; table 1). Table 2 shows procedural

ten patients did not have endovascular thrombectomy as

characteristics and complications.

there was no retrievable thrombus, whereas in the

bridging therapy group, no retrievable thrombus was

found in 15 patients who hence had no thrombectomy

(figure 1).

Baseline characteristics were similar in the two groups

(table 1). Overall, the median age of the patients was

68 years (IQR 61-78), and 166 (57%) of 293 were men. The

median NIHSS score was 15 in both groups, and the

median time from hospital admission to thrombolysis in

the bridging therapy group was 64 min (IQR 47-87). The

median time from randomisation to arterial puncture was

29 min (19-47) in the direct thrombectomy group and

42 min (29-59) in the bridging therapy group. Median

Articles

4b. Use the Select Tool:

To switch from the default Highlighting tool to the Select tool, click the middle icon above the full text (see red arrow in the top menu below).

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Nest Home

Literature Search

Screening

Tagging

Extraction

Risk of Bias

Study Inspector

Synthesis

Abstract

Full Text

Supplements

Related Reports

10

0

0

4

0

0

PubMed

Navigation

Tagging

Tag Recommendations

Comments (0)

History

Text Annotation

Direct thrombectomy

Bridging therapy

Effect size (95% CI)

p value

Primary efficacy outcome (ITT)

Functional independence: mRS 0-2 or return to baseline

80/146 (55%)

89/147 (61%)

Risk difference -0.051 (-0.160 to 0.059), adjusted OR 0.75 (0.45 to 1.24)

p=0.19 for non-inferiority; p=0.36 for superiority of bridging therapy

Primary efficacy outcome (PP)

Functional independence: mRS 0-2 or return to baseline

79/145 (54%)

88/143 (62%)

Risk difference -0.062 (-0.173 to 0.049), adjusted OR 0.69 (0.41 to 1.15)

p=0.25 for non-inferiority; p=0.36 for superiority of bridging therapy

Secondary outcomes (ITT)

mRS 0-1 or return to baseline

62/146 (42%)

71/147 (48%)

Adjusted OR 0.76 (0.46 to 1.24)

p=0.27

Score on mRS at 90 days

0

22/146 (15%)

30/147 (20%)

-

-

1

37/146 (25%)

40/147 (27%)

-

-

2

20/146 (14%)

18/147 (12%)

-

-

3

25/146 (17%)

19/147 (13%)

-

-

4

17/146 (12%)

13/147 (9%)

-

-

5

4/146 (3%)

5/147 (3%)

-

-

6

23/146 (16%)

24/147 (16%)

-

-

Score on ordinal analysis

2 (1-4)

2 (1-4)

-

-

Common adjusted OR 0.85 (0.56 to 1.27)

p=0.42

Thrombectomy mTICI score ≥3

127/143 (89%)

130/146 (89%)

Adjusted OR 0.54 (0.39 to 0.82)

p=0.66

NIHSS score within 72 h

4 (1-11), n=141

4 (1-11), n=142

Adjusted OR 0.73 (0.45 to 1.18)

p=0.20

Early neurological improvement*

84/141 (60%)

95/142 (67%)

Adjusted OR 0.73 (0.45 to 1.18)

p=0.20

Safety outcomes

Death

22/146 (15%)

24/147 (16%)

Adjusted OR 0.92 (0.46 to 1.84)

p=0.82

Symptomatic intracerebral haemorrhage

2/146 (1%)

3/147 (2%)

Adjusted OR 1.70 (0.22 to 13.04)

p=0.62

Any intracerebral haemorrhage

3/146 (2%)

3/147 (2%)

Adjusted OR 0.97 (0.56 to 1.70)

p=0.92

Data are n/N (%) or median (IQR). ITT=Intention to treat; mRS=modified Rankin Scale; PP=per protocol; OR=odds ratio; NIHSS=National Institutes of Health Stroke Scale; mTICI=modified Treatment in Cerebral Ischemia; *NIHSS reduction of 8 points or more, or reaching 0-1 at 3 days, adjusted for baseline NIHSS and age.

Table 3: Study outcomes

patients lost to follow-up. In the per-protocol analysis

times from arterial puncture to reperfusion were below

after patient crossover, 145 remained in the direct

60 min in both groups (55-5 min [26-0-88-5] in the direct

thrombectomy group and 143 in the bridging therapy

thrombectomy group and 44-5 min [27-0-70-0] in the

Create a box across the area you'd like to select for the tag (red arrow in the text section). Click in the left-hand corner of your area of interest and drag across the text or table. This selection will be automatically saved in the tag text box.

Selection / Area Annotation is best used on tables, figures, and images that are not amenable to exact

text quotation.

4c. Manually type out in Tag text box:

If you prefer to manually type the information from the text, you can do this by clicking your cursor in the tag text box and type what you'd like.

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Nest Home

Dashboard

Settings

Literature Search

1 / 1

Other Sources

Duplicate Review

Search Exploration

Query Builder

Screening

271 / 234

Configure Screening

Tagging

3 / 6

Configure Tagging

Extraction

6 / 6

Configure Extraction

Risk of Bias

6 / 6

Study Inspector

Synthesis

Manuscript Editor

Abstract Editor

Export

Abstract

Full Text

Supplements

Related Reports

10

0

4

0

0

PubMed

Navigation

Back

Skip

Complete

Tagging

Tag

Text

Select Tag

Between June 2, 2018, and July 8, 2021, 295 patients were randomly assigned to direct thrombectomy (n=148) or bridging therapy (n=147).

Apply Tag

Tag Recommendations

Comments (0)

History

Figure 2: Primary outcome overall and in prespecified Asian region subgroup

(A) Primary outcome by overall distribution by direct thrombectomy and bridging therapy (risk difference: -0.051, 95% CI -0.16 to 0.053, p=0.19). (B) Primary outcome distribution by direct thrombectomy and bridging groups in the Asian region prespecified subgroup (adjusted odds ratio 0.42, 95% CI 0.21 to 0.88, p=0.017). Horizontal stacked bar graphs show the primary outcome mRS distribution by direct thrombectomy and bridging therapy groups. Bars are labelled with proportions, mRS-modified Rankin Scale.

thrombectomy 14% 16% 13% 20% 1% 25%

bridging therapy 22% 22% 12% 7% 6% 22%

Proportion (%)

100

0

25

50

75

100

Role of the funding source

DIRECT-SAFE was an investigator-led clinical trial. The sponsor of the trial was the Florey Institute. The trial was supported by a grant from the Australian National Health and Medical Research Council programme, and Stryker USA. The trial was managed by Neuroscience Trials Australia. Database management and central data monitoring and verification was performed by Neuroscience Trials Australia and independent statistical analysis done by the Methods and Implementation Support for Clinical and Health Research Hub at the University of Melbourne, VIC, Australia. The regulatory sponsor and funding agents did not participate in the study design.

Results

Between June 2, 2018, and July 8, 2021, 295 patients were randomly assigned to direct thrombectomy (n=148) or bridging therapy (n=147). After two patients withdrew consent, 146 patients remained in the direct thrombectomy group and 147 patients in the bridging therapy group for intention-to-treat analysis, with no

generating stratum-specific risk differences with corresponding 95% CIs for each of the four strata (age <60 years vs 60 years or older by baseline NIHSS 0-15 or 16 and above) with subsequent pooling across strata using the Mantel-Haenszel method.

The proportions of mRS 0-2 or no change from baseline and death due to any cause were compared between the direct thrombectomy group and the standard bridging therapy group, adjusted for geographical region, age, and baseline NIHSS score using a logistic regression model. The proportions of participants with good angiographic reperfusion

120

www.thelancet.com Vol 400 July 9, 2022

Articles

	Direct thrombectomy	Bridging therapy	Effect size (95% CI)	p value
Primary efficacy outcome (ITT)				
Functional independence: mRS 0-2 or return to baseline	80/146 (55%)	89/147 (61%)	Risk difference: -0.051 (-0.160 to 0.053); adjusted OR 0.75 (0.42 to 1.34)	p=0.19 for non-inferiority; p=0.26 for superiority of bridging therapy

Manual text entry should be used whenever you want to associate customized text rather than quotation from the underlying article. **Warning:** manual entry will not maintain an exact location in the full text, so it may be difficult to find the exact contents of the article that support manually entered text excerpts.

Clear Annotations

If you need to redo your tag text annotation, you can either simply redo the action (Highlighting, Selecting, or Manually typing), or select "Clear Annotation" from the top of the Full Text:

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Nest Home

Dashboard

Settings

Literature Search

Other Sources

Duplicate Review

Search Exploration

Query Builder

Screening

271 / 274

Configure Screening

Tagging

3 / 6

Configure Tagging

Extraction

6 / 6

Configure Extraction

Risk of Bias

6 / 6

Study Inspector

Synthesis

Manuscript Editor

Abstract Editor

Export

Abstract

Full Text

Supplements

Related Reports

10

0

4

0

PubMed

3 / 6

Navigation

Back

Skip

Complete

Tagging

Tag

Text

Select Tag

DIRECT-SAFE was an investigator-led clinical trial. The sponsor of the trial was the Florey Institute. The trial was supported by a grant from the Australian National Health and Medical Research Council programme, and Stryker USA. The trial was managed by Neuroscience Trials Australia. Database management and central data monitoring and verification was performed by Neuroscience Trials Australia and independent statistical analysis done by the Methods and Implementation Support for Clinical and Health Research Hub at the University of Melbourne, VIC, Australia. The regulatory sponsor and funding agents did not participate in the study design.

Apply Tag

Tag Recommendations

Comments (0)

History

Thrombectomy

14%

35%

32%

22%

22%

45%

25%

Bridging therapy

22%

22%

18%

7%

8%

6%

21%

Proportion (%)

Figure 2: Primary outcome overall and in prespecified Asian region subgroup. (A) Primary outcome by overall distribution by direct thrombectomy and bridging therapy (risk difference -0.051; 95% CI -0.18 to 0.075; p=0.19). (B) Primary outcome distribution by direct thrombectomy and bridging groups in the Asian region prespecified subgroup (adjusted odds ratio 0.42; 95% CI 0.12 to 0.86; p=0.007). Horizontal stacked bar graphs show the primary outcome (mRS) distribution by direct thrombectomy and bridging therapy groups. Bars are labelled with proportions. mRS=modified Rankin Scale.

generating stratum-specific risk differences with corresponding 95% CIs for each of the four strata (age <60 years vs 60 years or older by baseline NIHSS 0-15 or 16 and above) with subsequent pooling across strata using the Mantel-Haenszel method.

The proportions of mRS 0-2 or no change from baseline and death due to any cause were compared between the direct thrombectomy group and the standard bridging therapy group, adjusted for geographical region, age, and baseline NIHSS score using a logistic regression model. The proportions of participants with good angiographic reperfusion

Role of the funding source
DIRECT-SAFE was an investigator-led clinical trial. The sponsor of the trial was the Florey Institute. The trial was supported by a grant from the Australian National Health and Medical Research Council programme, and Stryker USA. The trial was managed by Neuroscience Trials Australia. Database management and central data monitoring and verification was performed by Neuroscience Trials Australia and independent statistical analysis done by the Methods and Implementation Support for Clinical and Health Research Hub at the University of Melbourne, VIC, Australia. The regulatory sponsor and funding agents did not participate in the study design.

Results
Between June 2, 2018, and July 8, 2021, 295 patients were randomly assigned to direct thrombectomy (n=148) or bridging therapy (n=147). After two patients withdrew consent, 146 patients remained in the direct thrombectomy group and 147 patients in the bridging therapy group for intention-to-treat analysis, with no

120

www.thelancet.com Vol 400 July 9, 2022

Articles

	Direct thrombectomy	Bridging therapy	Effect size (95% CI)	p value
Primary efficacy outcome (ITT)				
Functional independence: mRS 0-2 vs return to baseline	80/146 (55%)	89/147 (61%)	Risk difference: -0.051 (-0.160 to 0.059); adjusted OR 0.75 (0.40 to 1.24)	p=0.19 for non-inferiority; p=0.25 for superiority of bridging therapy

This will remove all tag text; next, choose the tag text type you would like to use, and redo the relevant Highlight, Selection, or Manual text entry.

Q: Why not leave the annotation / tag text blank?

A: It is possible to apply tags without filling in the tag text. However, doing so will mean that the only evidence that the tag is applicable to that specific study will be the fact that it was applied, and those who view your Qualitative Synthesis will have no context. If you fill in text content, you provide specific evidence of that tag's applicability as well as presenting the specific information from that study to viewers of Qualitative Synthesis.

5. Click "Apply Tag"

Once you have the content of interest into the tag text box, make sure that you have selected the relevant tag from the drop-down menu (red box). Once you have confirmed that both the Tag and the Tag Text Content are correct, click "Apply Tag."

Nested Knowledge - <https://wiki.nested-knowledge.com/>

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

Navigation: Skip Complete

Tagging: Tag Text mRS 0-2 Apply Tag

Tag Recommendations: Comments (0) History

	Direct thrombectomy	Bridging therapy	Effect size (95% CI)	p value
Primary efficacy outcome (ITT)				
Functional independence: mRS 0-2 or return to baseline	80/145 (55%)	89/147 (61%)	Risk difference: -0.053 (-0.160 to 0.053) adjusted OR 0.75 (0.45 to 1.24)	p=0.19 for non-inferiority; p=0.26 for superiority of bridging therapy
Primary efficacy outcome (PP)				
Functional independence: mRS 0-2 or return to baseline	79/145 (54%)	88/143 (62%)	Risk difference: -0.062 (-0.173 to 0.049) adjusted OR 0.69 (0.41 to 1.15)	p=0.25 for non-inferiority; p=0.16 for superiority of bridging therapy
Secondary outcomes (ITT)				
mRS 0-1 or return to baseline	62/145 (43%)	71/147 (48%)	Adjusted OR 0.76 (0.46 to 1.24)	p=0.27
Score on mRS at 90 days				
0	22/145 (15%)	30/147 (20%)	-	-
1	37/145 (26%)	40/147 (27%)	-	-
2	20/145 (14%)	19/147 (13%)	-	-
3	25/145 (17%)	19/147 (13%)	-	-
4	17/145 (12%)	13/147 (9%)	-	-
5	4/145 (3%)	5/147 (3%)	-	-
6	2/145 (1%)	2/147 (1%)	-	-
Score on ordinal analysis	7 (3-9)	7 (3-9)	Common adjusted OR 0.85 (0.56 to 1.27)	p=0.42
Thrombectomy mRS score 2b-3	122/143 (85%)	130/146 (89%)	Adjusted OR 0.84 (0.39 to 1.82)	p=0.66
NIHSS score within 72 h	4 (1-11), n=141	4 (1-11), n=142	-	-
Early neurological improvement*	84/141 (60%)	95/142 (67%)	Adjusted OR 0.73 (0.45 to 1.18)	p=0.20
Safety outcomes				
Death	22/145 (15%)	24/147 (16%)	Adjusted OR 0.92 (0.45 to 1.84)	p=0.82
Symptomatic intracerebral haemorrhage	2/145 (1%)	1/147 (1%)	Adjusted OR 1.70 (0.22 to 13.04)	p=0.63, Fisher's exact test
Any intracerebral haemorrhage	22/145 (15%)	22/147 (15%)	Adjusted OR 0.97 (0.56 to 1.70)	p=0.92

Data are n/N (%) or median (IQR). ITT: intention to treat; mRS: modified Rankin Scale; PP: per protocol; OR: odds ratio; NIHSS: National Institutes of Health Stroke Scale; mRS: modified Rankin Scale; mRS: modified Rankin Scale; mRS: modified Rankin Scale. *NIHSS reduction of 8 points or more, or reaching 0-1 at 3 days, adjusted for baseline NIHSS and age.

Table 3: Study outcomes

Note: Anytime there is a module box with the adjustable icon, you can drag to adjust the width of the box depending on your preference.

Navigation: Skip Complete

Tagging: Tag Contents Select Tag Enter Text Apply Tag

Tag Recommendations: Comments (0) History

Original Research

Health complaints and use of medicines among adolescents in Malta

Rita DARMANIN ELLUL, Maria CORDINA, Anton BUHAGIAR, Anthony FENECH, Janet MIFSUD.

Received (first version): 17-Mar-2008 Accepted: 16-Aug-2008

ABSTRACT

Objective: To investigate self-reported health complaints and the use of medicines among adolescents in Malta.

Methods: A self-administered questionnaire was used to survey self-reported health complaints, the adolescents that will integrate information about the proper use of medicines.

Keywords: Adolescent. Drug Utilization. Malta.


Tagging Supplemental Materials

If the study you are tagging has supplemental materials in pdf format, you may also apply tags to these texts. The functionality is the same as tagging full texts: automatic copy of text to clipboard and text box, text highlighting and area selection allowing immediate direction to the excerpt when the tag is selected etc.

Note: Tag Recommendations are currently unavailable for supplemental pdfs.

AbstractFull TextSupplementsRelated ReportsCT.gov

Author Manuscript

**HHS Public Access**
Author manuscript
Indian J Pediatr. Author manuscript; available in PMC 2016 June 01.

Published in final edited form as:
Indian J Pediatr. 2012 August ; 79(8): 1062–1068. doi:10.1007/s12098-012-0765-1.

Vitamin D in Chronic Kidney Disease

Yahn-Yir Chau and
Taipei Cathay General Hospital, Taipei, Taiwan

Juhi Kumar
Weill Cornell Medical College, New York, NY, USA

Navigation

BackSkipComplete

Questions (0/59)

Tagging

TagContents

Sex

Enter Text

Apply Tag

History

Tags with Table Contents

Similarly to tags with text contents, you select the dropdown to find the tag of choice. When selected, the table you created will be shown and you can input text into any of the rows. When you are satisfied, click “Apply Tag.”

Tagging: Thrombectomy alone vs. Thrombectomy plus thrombolysis

AbstractFull TextSupplementsRelated Reports13541484PubMed

Nest Home

Dashboard

Settings

Literature Search1/1

Other Sources

Duplicate Review

Search Exploration

Screening271/293

Configure Screening

Tagging3/6

Configure Tagging

Extraction6/6

Configure Extraction

Risk of Bias6/6

Study Inspector

Synthesis

Manuscript Editor

Abstract Editor

Export

Table 1. Demographic and Clinical Characteristics of the Patients at Baseline.*

Characteristic	EVT Alone (N=273)	Alteplase Followed by EVT (N=266)
Median age (IQR) — yr	72 (62–80)	69 (61–77)
Male sex — no. (%)	161 (59.0)	144 (54.1)
Median NIHSS score (IQR)†	16 (10–20)	16 (10–20)
Medical history		
Ischemic stroke — no. (%)	47 (17.2)	44 (16.5)
Atrial fibrillation — no. (%)	86 (31.5)	63 (23.7)
Diabetes mellitus — no. (%)	40 (14.7)	50 (18.8)
Hypertension — no./total no. (%)	121/273 (44.3)	139/265 (52.5)
Prestroke score on the modified Rankin scale — no./total no. (%)‡		
0	189/272 (69.5)	185/266 (69.5)
1	51/272 (18.8)	49/266 (18.4)
2	24/272 (8.8)	25/266 (9.4)
≥3	8/272 (2.9)	7/266 (2.6)
Median systolic blood pressure (IQR) — mm Hg§	150 (135–167)	150 (130–169)
Median glucose level (IQR) — mmol/liter¶	6.6 (5.8–7.6)	6.8 (5.9–8.5)
Median ASPECTS (IQR)‡	9 (8–10)	9 (8–10)
Location of intracranial occlusion — no./total no. (%)**		
Intracranial ICA	4/272 (1.5)	0/266
Terminal ICA	64/272 (23.5)	50/266 (18.8)
M1	156/272 (57.4)	174/266 (65.4)

Tagging

TagContents

Sex

MaleFemalePrefer not to say

205333

Apply Tag

Tag Recommendations

Comments (0)

History

Highlighting pdfs does not automatically input the text into the box unlike tags with text contents only. However, it will remember any text highlighted or selected in the pdf and auto-scroll to it when the tag is selected again.


Note: If you are entering numerical data into tables, no automated statistics are generated. This is only done in the Meta-Analytical Extraction module.

To alter the columns in the table for this tag, either click on the column header in the Tagging module itself, or head back to Configure Tagging. [Learn more about tag tables here.](#)

Add New Tags on the Fly

When you find a term that you want to add to the Tag Hierarchy, you can either add it on the Configure Tagging page, or add it 'on the fly' without leaving the page.

To add a tag on the fly, type the title of your new tag into the “Select Tag” box, and click “Add Option” that appears at the top of the drop-down list of tags.



The image shows a modal window titled "NEW TAG NAME". At the top, there is a header bar with a dropdown arrow, the number "0", and an upward arrow. Below the header, there is a list of parent tags for selection. The first item is "Add Option: NEW TAG NAME" in a dark blue bar. The other items are in light blue bars and include: "Patient Characteristics", "Rescue therapy applied?", "Aspiration", "IVT", "Inclusion Criteria for Occlusion Location", "Posterior Cerebral Artery (PCA)", "Anterior Cerebral Artery (ACA)", "Middle Cerebral Artery", "Basilar artery", "Internal carotid artery (ICA)", "Parent Artery Location", "Last Known Well", "Presentation", "NIHSS (mean) Baseline", "ASPECTS", and "NIHSS (median) Baseline". A vertical scrollbar is on the right side of the list.

In the modal that appears, confirm the tag name, add a description (optional), and as relevant, identify the new tag's Parent Tag. Once created, you will now be able to find the new Tag on the drop-down list.

Note: Only tags with text contents can be created on the fly. To toggle on table contents, edit the tag in Configure Tagging.

From:

<https://wiki.nested-knowledge.com/> - **Nested Knowledge**

Permanent link:

<https://wiki.nested-knowledge.com/doku.php?id=wiki:autolit:tagging:tag&rev=1698927407>

Last update: **2023/11/02 12:16**