

Nest Protocol


Purpose

The Nest Protocol outlines the goal, scope, and key tasks of your systematic literature review or meta-analysis. In it, the project leadership should provide critical background information, the context of the review, and primary and secondary endpoints.

Editing the Protocol

While the Protocol can be viewed from Nest Home, it is only editable from [Nest Settings](#).

1. Navigate to Settings



2. Select Protocol Editor

Administrator Settings: Basilar Artery - thrombectomy vs. thrombolysis

Synthesis

Choose outputs to display on Synthesis:

Checking these boxes will show the respective outputs on Synthesis for this nest

☒ Qualitative Synthesis

☒ Quantitative Synthesis

☒ Manuscript

Make this nest public:

Checking this box will allow anyone on the internet to search for and see your nest's Synthesis. This does not grant the public editing access via AutoLit

☒ Public Nest

Static Manuscript

Upload a PDF-based manuscript for this nest. Displayed only if the editable Manuscript is empty.

Upload .pdf

Edit Nest Description:

Write a brief, public description of your nest, as it should appear on Synthesis

Placeholder

Saved

Protocol Editor

The Protocol Editor allows you to create and maintain Nest-specific documentation linked to particular parts of the AutoLit workflow.

Write the Protocol

Add headers, bullet points, and bold font. There is no one-size-fits all formula for a protocol! The goal of a 'best practices' protocol should be: enable someone unfamiliar with the project to carry out any of the Search, Screening, Tagging, Extraction, or interpretive tasks in your project in the same way that you, as project lead, would do.

Show Table of Contents

Protocol

Description

Objective

To determine the importance of screening for Dihydropyrimidine Dehydrogenase (DPD) deficiency in patients undergoing 5-FU based chemotherapy

Scope

Clinical trials, retrospective studies or prospective studies evaluating outcomes related to toxicities in patients undergoing 5-FU based chemotherapy with DPD deficiency versus with no DPD deficiency.

Primary Outcomes

- Risk of Severe adverse effects (*grade 2 or above*) in population with dpd deficiency
- Allele frequency
- Prevalence of dpd deficiency in patients undergoing 5\ -fu based chemotherapy for various cancers

Secondary Outcomes

- Effect of combination therapies
- Type of combination drug attributable to the serious adverse effect
- Type of the serious adverse effects with the therapy
- Percentage of DPD negative patients with adverse effects
- Percentage of DPD positive patients with no adverse effects

Protocol Best Practices

To see general practice recommendations on protocol drafting (and a generic template), see [Protocol Best Practices](#).

From:

<https://wiki.nested-knowledge.com/> - **Nested Knowledge**

Permanent link:

<https://wiki.nested-knowledge.com/doku.php?id=wiki:autolit:admin:protocol&rev=1653325988>

Last update: **2022/05/23 17:13**